



**University of  
Zurich<sup>UZH</sup>**

**Zurich Open Repository and  
Archive**

University of Zurich  
University Library  
Strickhofstrasse 39  
CH-8057 Zurich  
[www.zora.uzh.ch](http://www.zora.uzh.ch)

---

Year: 2018

---

## **Prediction of ROP Treatment and Evaluation of Screening Criteria in VLBW Infants-a Population Based Analysis**

Gerull, Roland ; Brauer, Viviane ; Bassler, Dirk ; Laubscher, Bernard ; Pfister, Riccardo E ; Nelle, Mathias ; Müller, Beatrice ; Roth-Kleiner, Matthias ; Gerth-Kahlert, Christina ; Adams, Mark ; Swiss Neonatal Network ; Follow-up Group

**Abstract:** BACKGROUND The incidence of retinopathy of prematurity (ROP) and ROP screening criteria differ between countries. We assessed whether ROP screening could be reduced based on the local ROP incidence. METHODS Observational cohort study of infants born in Switzerland between 2006 and 2015 <32 0/7 weeks. Chronological and postmenstrual ages at ROP treatment were analyzed. A model to identify ROP treatment on patients born between 2006 and 2012 (training set) was developed and tested on patients born between 2013 and 2015 (validation set). RESULTS Of 7817 live-born infants, 1098 died within the first 5 weeks of life. The remaining 6719 infants were included into analysis. All patients requiring ROP treatment would have been identified if screening had been performed before reaching 60 days of life or 37 3/7 weeks postmenstrual age, whichever came first. The training and validation sets included 4522 and 2197 preterm infants encompassing 56 and 20 patients receiving ROP treatment, respectively. All patients would have required screening to reach 100% sensitivity. To reach a sensitivity of 95.0% and a specificity of 87.6%, we predicted a reduction in 13.2% of patients requiring screening (c-statistic = 0.916). CONCLUSIONS A substantial reduction of infants requiring screening seems possible, but necessitates prospective testing of new screening criteria.

DOI: <https://doi.org/10.1038/s41390-018-0128-3>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-156862>

Journal Article


Accepted Version

Originally published at:

Gerull, Roland; Brauer, Viviane; Bassler, Dirk; Laubscher, Bernard; Pfister, Riccardo E; Nelle, Mathias; Müller, Beatrice; Roth-Kleiner, Matthias; Gerth-Kahlert, Christina; Adams, Mark; Swiss Neonatal Network; Follow-up Group (2018). Prediction of ROP Treatment and Evaluation of Screening Criteria in VLBW Infants-a Population Based Analysis. *Pediatric Research*, 84(5):632-638.

DOI: <https://doi.org/10.1038/s41390-018-0128-3>

Author Version: Published ahead of online first



## Prediction of ROP Treatment and Evaluation of Screening Criteria in VLBW Infants—a Population Based Analysis

Roland Gerull, Viviane Brauer, Dirk Bassler, Bernard Laubscher, Riccardo E Pfister, Mathias Nelle, Beatrice Müller, Matthias Roth-Kleiner, Christina Gerth-Kahlert, Mark Adams

**Cite this article as:** Roland Gerull, Viviane Brauer, Dirk Bassler, Bernard Laubscher, Riccardo E Pfister, Mathias Nelle, Beatrice Müller, Matthias Roth-Kleiner, Christina Gerth-Kahlert and Mark Adams, Prediction of ROP Treatment and Evaluation of Screening Criteria in VLBW Infants—a Population Based Analysis, *Pediatric Research* \_\_\_\_\_ doi:10.1038/s41390-018-0128-3

This is a PDF file of an unedited peer-reviewed manuscript that has been accepted for publication. Springer Nature are providing this early version of the manuscript as a service to our customers. The manuscript will undergo copyediting, typesetting and a proof review before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers apply.

**Title:**

Prediction of ROP Treatment and Evaluation of Screening Criteria in VLBW Infants – a Population Based Analysis

**Running title**

ROP Prediction in Switzerland

**Author listing**

Roland Gerull (1, 2)

Viviane Brauer (1)

Dirk Bassler (3)

Bernard Laubscher (4)

Riccardo E Pfister (5)

Mathias Nelle (1)

Beatrice Müller (6)

Matthias Roth-Kleiner (7)

Christina Gerth-Kahlert (8)

Mark Adams (3)

on behalf of the Swiss Neonatal Network & Follow-up Group

(1) Inselspital Bern, Neonatology, University of Berne, 3008 Berne, Switzerland

(2) University of Basel, Children's Hospital UKBB, Department of Neonatology, 4056 Basel, Switzerland

(3) University Hospital Zurich, Department of Neonatology, 8091 Zurich, Switzerland

(4) Hôpital Neuchâtelois, Pediatrics, 2000 Neuchâtel, Switzerland

(5) University Hospital Geneva, Neonatology, 1211 Geneva, Switzerland

(6) Ostschweizer Kinderspital St. Gallen, Intensive Care and Neonatology, 9006 St. Gallen, Switzerland

(7) University Hospital and University of Lausanne, Clinic of Neonatology, 1011 Lausanne, Switzerland

(8) University Hospital Zurich, Department of Ophthalmology, 8091 Zurich, Switzerland

**Corresponding Author**

Roland Gerull

University of Basel, Children's Hospital UKBB, Department of Neonatology, Spitalstrasse 33, 4056 Basel, Switzerland

roland.gerull@ukbb.ch

Phone: 061 704 2307

Fax: 061 265 9568

**Statement of financial support**

No sources of external funding.

**Disclosure statement**

The authors declare no conflict of interest.

**Category of study**

Population study

**Abstract**

**Background:** The incidence of retinopathy of prematurity (ROP) and ROP screening criteria differ between countries. We assessed whether ROP screening could be reduced based on the local ROP incidence.

**Methods:** Observational cohort study of infants born in Switzerland between 2006 and 2015 <32 0/7 weeks. Chronological and postmenstrual ages at ROP treatment were analyzed. A model to identify ROP treatment on patients born 2006-2012 (training set) was developed and tested on patients born 2013-2015 (validation set).

**Results:** Of 7817 live-born infants, 1098 died within the first five weeks of life. The remaining 6719 infants were included into analysis. All patients requiring ROP treatment would have been identified if screening had been performed before reaching 60 days of life or 37 3/7 weeks postmenstrual age, whichever came first.

The training and validation sets included 4522 and 2197 preterm infants encompassing 56 and 20 patients receiving ROP treatment, respectively. All patients would have required screening to reach 100% sensitivity. To reach a sensitivity of 95.0% and a specificity of 87.6% we predicted a reduction to 13.2% of patients requiring screening (c-statistic = 0.916).

**Conclusions:** A substantial reduction of infants requiring screening seems possible but necessitates prospective testing of new screening criteria.

## Introduction

Retinopathy of prematurity (ROP) is a severe complication of preterm birth and may lead to severe visual impairment or even blindness. It is a two stage developmental vascular proliferative disorder resulting from a mismatch between oxygen demand and oxygen supply within the retina (1, 2). By now a large variety of risk factors for the development of ROP have been described, but it is not clear which risk factors are truly independent and which factors are rather associations (3).

In developed countries the incidence of ROP in preterm infants below 28 weeks of gestational age (GA) ranges widely between 25 and 91%, with treatment rates varying between 2% and 30% (4). Criteria for ROP screening differ between countries and, in developed countries, usually include patients below 31-32 weeks of gestational age (GA) and a birthweight (BW) below 1250-1500g. All recommendations extend ROP screening to patients above given limits for gestational age and birth weight if additional risk factors were present. These factors are not clearly defined but include prolonged oxygen therapy (Argentina(5)), respiratory problems or sepsis (Brazil (6)), “infants believed to be at high risk for ROP” (Canada (7)), or “unstable clinical course with cardiorespiratory support and are believed to be at high risk of ROP” (USA(8)).

However, these criteria seem to miss patients potentially requiring treatment in developing countries (9, 10). Screening inclusion criteria for ROP need to reach a high sensitivity to avoid missing patients that potentially could suffer severe consequences such as blindness. When prevalence is small, high sensitivity leads to low specificity with many patients screened that ultimately do not require treatment. The lower the prevalence for infants requiring ROP treatment, the more crucial it is to optimize screening criteria in order to reduce work load in the NICUs, financial expenses and, most importantly, avoidable stress for preterm infants. This is an issue in particular in Switzerland with its very low prevalence of

ROP treatment with 4.3% compared to other countries like Canada (10.4%), United Kingdom (8.5%) or Australia & New Zealand with 7.3% in preterm infants below 28 weeks (4, 11).

The aim of this study was to assess if screening criteria for ROP can be optimized in very low birth weight (VLBW, birth weight below 1500g) infants in Switzerland. All VLBW infants live-born between 2006 and 2015 in Switzerland were included in this study.

## Patients and Methods

This study is based on a retrospective analysis of patients who were born between 2006 and 2015 and were registered at the national registry of very preterm infants in Switzerland (SwissNeoNet, SNN) of the Swiss Society of Neonatology. The network prospectively collects perinatal and follow-up data of live born infants with a gestational age between 22 0/7 weeks and less than 32 0/7 weeks or a birth weight of less than 1501g. All 9 Swiss perinatal centers, 5 step-down units and 16 neuro-/developmental pediatric units participated. Routine comparison with the Swiss Federal Statistical Office reveals 96% population coverage. This study includes patients with a gestational age below 32 0/7 weeks and focused on data from the perinatal dataset plus the information of any ROP treatment after primary discharge home from the 18 to 24 months follow-up assessment. Infants with a gestational age as of 32 0/7 and a birth weight below 1501g were excluded because a cohort with both gestational age and birth weight as cutoff cannot include gestational age as main predictor. None of these infants were treated for ROP. We excluded all infants that died before reaching 5 weeks of age as they never would have received a ROP screening.

Screening criteria for ROP differed slightly between the individual centers. ROP screening criteria included a birth weight of <1500g in eight of nine Swiss hospitals with tertiary neonatal care. The remaining center screened only patients with a birth weight below 1250g. In seven hospitals preterm infants with a gestational age of less than 32 0/7 weeks were screened, while in the remaining two centers the cutoff was at 30 0/7 weeks. One center started screening at a postnatal age of more than seven weeks; all other centers started ROP screening between 4 and 5 weeks. Administration of supplemental oxygen for  $\geq 3$  days or unstable clinical course was defined as additional screening criterion in most hospitals. The final model contained the following 5 risk factors: GA, birth weight z-score, CPAP > 3 days, multiple birth and surfactant. The parameters rejected were: male sex, caesarean section, any antenatal steroids, delivery room intubation and mean weight gain per week hospital stay, any mechanical ventilation, supplemental oxygen > 3 days. The date of first ROP screening differed between 6 and 8 weeks postnatally. ROP was assessed by indirect ophthalmoscopy by the attending ophthalmologists. In Switzerland, all ROP treatments are performed by one of five ophthalmological clinics.

## Definitions

Stages of ROP were categorized according to the international classification of retinopathy of prematurity (12). Severe ROP (sROP) was defined as ROP stages 3-5. Data on retinal zones and development of plus disease were not available from the database. ROP was stratified as treated ROP when any form of therapy (laser coagulation or intravitreal anti-VEGF) was performed according to current recommendations.

Patients were classified as small for gestational age (SGA) when the birth weight was below the 10<sup>th</sup> percentile based on the growth curves by Voigt et al. (13). Length of support with

supplemental oxygen, CPAP and mechanical ventilation was measured as number of days during hospitalization with more than 12 hours of the different form of respiratory support, respectively. We defined bronchopulmonary dysplasia (BPD) as an oxygen requirement at 36 weeks postmenstrual age according to the NICHD consensus conference paper (14), NEC as clinical signs (abdominal distension, bilious aspirates and/or bloody stools) confirmed by radiographically visible intramural gas or at laparotomy (Bell stages 2 and 3) (15), and antenatal steroid use as any administration prior to birth, regardless of the time interval. Growth per week was calculated as weight at discharge or death minus weight at birth divided by the amount of weeks from birth until first discharge home or until death. Sepsis was defined as clear clinical evidence of infection plus at least one positive blood culture (including coagulase negative staphylococci and fungal pathogens). Severe intraventricular hemorrhage (sIVH) was based on the most severe ultrasound result during hospital stay reaching grade 3 to 4 of the classification defined by Papile et al. (16).

## Ethics

Data collection and evaluation for this study were approved by the Swiss Federal Commission for Privacy Protection in Medical Research and the Swiss ethical review boards (KEK-ZH-Nr. 2014-0551 and KEK-ZH-Nr. 2014-0552). The patients' representatives were informed about the use of data for research.

## Statistical methods

Based on risk factors of patients born 2006-2012 (training set), a mathematical model using logistic regression for the outcome ROP intervention was performed using a step-wise elimination process while ensuring that variable elimination did not significantly change the model. An elimination of a variable was accepted only if the model with the variable was not



significantly different to the model without the variable according to Wald ( $p < 0.10$ ) (17). The final model contained the following 7 risk factors: GA, days of supplemental oxygen, days of CPAP, days of mechanical ventilation, birth weight z-score, surfactant, and multiple birth. The parameters rejected were: male sex, caesarean section, any antenatal steroids, delivery room intubation and mean weight gain per week hospital stay. To test validity of this model to predict preterm infants requiring ROP intervention it was applied on patients born 2013-2015 (validation set). Sensitivity and specificity were analyzed as well as ROC curves and predictive c-statistics. Model building and validation was based on five-fold imputed data using multivariate imputation by chained equations based on random forest. Three of the included parameters are not available at birth: duration of supplemental oxygen, CPAP and mechanical ventilation. We included a sensitivity analysis using these parameters as categorical data; whether or not they had longer than median length of respiratory support (supplemental Oxygen  $> 3$  days, CPAP  $> 3$  days, mechanical ventilation  $> 0$  days). Cochrane-Armitage test for trend was used to test the decrease of mortality over time. Results with a p-value of less than 0.05 were considered significant. Analyses were performed with SPSS Version 21 (IBM Corporation, Armonk, NY) and R 3.4 (r-project for statistical computing; [www.r-project.org](http://www.r-project.org)).

The following prediction model formula was applied:

$$p(y = 1 | x_1, \dots, x_p) = \frac{\exp(b_0 + \text{Sum}(b_k * x_k))}{1 + \exp(b_0 + \text{Sum}(b_k * x_k))}$$

$b_0$  = intercept;  $b_k$  = maximum likelihood estimate for chosen risk factors;  $x_k$  = risk factor values of individual patient.

## Results

Record completeness of all infants according to comparison with Swiss Federal Statistical Office was 96%. Of the 7817 patients registered in the SNN database, information on ROP treatment was missing for 1116 patients of whom we retrospectively collected and completed 942 cases, leading to a data completeness of 97.8%. After exclusion of patients who died in the delivery room or during the first 5 weeks of life in hospital, 6719 infants remained for analyses.

Characteristics of included patients eligible to ROP screening are shown in table 1. Analyses showed that the majority of patients requiring ROP therapy were born with a gestational age below 28 weeks. ROP was treated in only 13, 6 and 3 patients at a gestational age of  $\geq 27$  0/7,  $\geq 28$  0/7 and  $\geq 29$  0/7 weeks, respectively (Figure 1). Patient characteristics of these patients are shown in table 2. The mortality at five weeks of age, after which ROP screening usually begins, significantly declined between 2006 and 2015 ( $p = 0.03$ ) raising the proportion of infants at risk for developing ROP over the years.

Infants were treated for ROP at a GA of  $38.17 \pm 3.46$  weeks, range 32.85 – 56.42 weeks. Only two patients were treated with a lower gestational age than 34 0/7 weeks and in three patients ROP therapy was performed after 44 weeks GA. The chronological age at ROP treatment was  $88 \pm 24$  days with a range from 41 to 215 days with one patient being treated before 60 days of life. The combination of GA and chronological age showed that all patients would have been identified prior to ROP treatment when screening had begun at 60 days of life or a GA of 37.42 weeks whichever is reached first.

GA-stratified analysis revealed that missing data were randomly distributed for most of the parameters. Antenatal steroids had a higher rate of missing data at 23 weeks gestational age (17.2%) and 24 weeks (5.4%), respectively, ROP intervention had a higher rate of missing data in infants with a gestational age of 31 weeks (5.3%). Logistic regression analysis was based on five-fold imputed data to compensate for missing values in antenatal steroids (2.7%), ROP intervention (2.6%) and to a minor degree in other parameters. Without data imputation, all cases with missing information for either the primary outcome (ROP intervention) or one of the predictors would have been eliminated from the model prior to model building. This would have led to the higher rate of exclusion of data on either end of the gestational age range.

The training set (patients born 2006-2012) had a patient population of 4522 preterm infants <32 weeks GA, including 56 patients with treated ROP. The validation set (patients born 2013-2015) included 2197 patients with 20 ROP therapies. Table 3 shows values, which resulted from analyses of our patient characteristics and were included in the final model.

Depending on the cutoff-point, the logistic regression model allows predicting the number of false negative cases, i.e. cases for which a ROP intervention was performed during 2013-2015 but which was not detected by the model. Number of infants needed to test, true positives, false negatives, sensitivity and specificity for each cutoff value are displayed in table 4.

Results show that all patients needed to be screened to reach a sensitivity of 100%. However, to reach a sensitivity of 95.0% (one patient false negative) and specificity of 87.6% the model predicted a reduction in the number of screened patients to 13.2%. This reflects a high predictive c-statistics value of 0.916 (Figure 2). The undetected patient was born at 30+1 weeks with congenital nephroblastoma and therefore probably with a different

pathophysiology than the average patient requiring ROP treatment towards which the prediction model was fit.

The analysis, using supplemental oxygen, duration of CPAP and duration of mechanical ventilation as categorical variables showed almost identical results. The ROC showed an AUC of 0.909. Only 16.7% of patients needed to be screened for a sensitivity of 95% and a specificity of 84.0% (details are presented in the supplemental material).

## Discussion

This study investigated ways to improve ROP screening criteria in Switzerland. Based on the low prevalence of ROP therapy during a period of ten years of 1.2% in children born below 32 weeks GA and less than 0.1% in infants born between 28 and 32 weeks, a logistic regression model was developed using known risk factors for ROP. The prediction model showed that all patients would have to be screened to guarantee a sensitivity of 100%. However, only 13% of patients needed to be screened for a sensitivity of 95.0%, missing one patient requiring ROP treatment.

All screening programs aim for a high sensitivity, since no patients with the specific condition should be missed. This is particularly true for ROP, where an unidentified infant can suffer dramatic consequences such as severe visual impairment or even blindness, underlining the importance of achieving 100% sensitivity. On the other hand high sensitivity in turn leads to a low specificity. This exposes a high number of patients to minor consequences, which in ROP screening means exposure to pain, stress, increased work load and higher financial expenses. Therefore, screening criteria should identify patients at risk for ROP and limit screening to these patients as far as safely possible. Furthermore, the latest safe time to begin screening should be identified.

Screening criteria for ROP usually include birth weight and gestational age. In developed countries most recommendations state that patients with birth weight of less than 1250-1500g or a gestational age of less than 30-32 weeks should be screened for ROP(18).

In developing countries recommendations frequently include patients with higher gestational age or higher birth weight (18). Several studies have shown that the incidence of ROP is higher in these countries and that application of criteria as defined by the American Academy of Pediatrics (AAP) (8) or the Royal College of Pediatrics and Child Health(19) misses patients needing ROP treatment (9, 10). This discrepancy confirms the absence of uniform screening criteria. Instead, we suggest that criteria could be improved if based on local risk factors and local incidence of ROP.

Our prediction model including data imputation showed that only 13% of patients needed to be screened for ROP to reach a sensitivity of 95.0% missing one patient. This raises the question if there were additional risk factors in this patient and if this patient should have been screened for the above mentioned unclearly defined criteria. These criteria could not be included in the prediction model since databases rely on variables with clear definitions and therefore cannot include parameters such as “believed to be at risk”.

The only patient missed with this prediction model suffered from a nephroblastoma. Nephroblastoma is a well vascularized tumor which is dependent on vasculogenesis for growth and which frequently expresses high doses of vascular growth factors, such as VEGF(20). It is well known, that VEGF plays a major role in the development of ROP in the second phase of relative hypoxia of the retina (2). Thus, it is very likely that the development of ROP in this patient was intensified by the growth factors produced by the tumor and not only by the usual pathophysiologic mechanisms occurring during ROP development in

preterm infants. Therefore, this patient had additional specific risk factors that make ROP screening necessary, independent of gestational age or birth weight.

Altogether, these considerations indicate that there is a large potential to reduce ROP the indications for screening in Switzerland. Our model applies data available at patient discharge (duration of supplemental oxygen, CPAP and mechanical ventilation) and is therefore not fit for direct prospective implementation at screening age. To assess if a similar model can be achieved with variables that are available before first screening is performed, we included a second sensitivity analysis using modes of respiratory support as categorical variables. The results were almost identical to the first analysis, proving large potential to reduce ROP screening. However, a prospective study first needs to be performed based on these screening criteria and augmented by a list of diagnosis for which ROP screening should be performed (independent of gestational age or birth weight in very preterm infants). This should enable the selection of new screening criteria matching the local situation.

Several studies assessed prediction models for ROP development. A recent publication summarized publications and methods which have been evaluated (21). Of these, the WINROP algorithm has been studied most extensively. WINROP was developed in Sweden and is based on postnatal weight gain, either alone (22) or in combination with serum levels of Insulin like growth factor 1 (IGF1) (23). The algorithm has mostly been tested retrospectively and was used in different populations. While prediction has shown to be excellent in one Swedish publication (24), application of the algorithm did not reach a sensitivity of 100% in all other populations with values mostly around 90% (25-28). Furthermore, these retrospective analyses included relatively small sample sizes of approximately 600 patients. This shows that a screening solely based on WINROP would fail to detect several patients developing severe ROP.

An extensive analysis of all Danish patients treated for ROP in the years 2002-2006 compared screening criteria combining gestational age at delivery and birth weight limits and new risk-based criteria were compared with regards to their effectiveness (3). Results showed that a reduction of 17.4% of screened patients allowed the detection of all patients treated for ROP in the observation period and might lead to one missed treatment-demanding ROP every 11 years and one case of blindness every 18 years. This implies that a potential for reduced ROP screening exists in Denmark. However, it also mirrors the difficulties of optimizing screening criteria with possibilities to avoid screening in a large group of preterm infants on the one hand but increasing risk to miss patients requiring ROP treatment on the other hand.

Besides the question who needs to be screened for ROP it is also a matter of debate when screening should be started. The AAP recommends that the first ROP screening should be performed at a chronological age of 4 weeks but not before a postmenstrual age of 31 weeks is reached(8). These limits are considerably earlier than results of our study, where all patients would have been detected prior to therapy with limits of a chronological age of 60 days or postmenstrual age of 37 weeks. This shows that the development of ROP differs between populations not only concerning of incidence but also in terms chronologic progression. Starting ROP screening in Switzerland at a later chronologic and/or postmenstrual age than at the dates recommended by the AAP seems possible.

Our study has several strengths and limitations. Of note is that different screening criteria were used across the neonatal centers, in particular some centers did not screen patients between 30 and 32 weeks GA. However, these units reduced screening because long periods of ROP-free infants above these criteria were documented. No extra cases of ROP were found at 2 years of age that were missed. Our study was based on a whole population in which a low

incidence of treated ROP was reported. This resulted in only 76 patients with treated ROP, which made the development, the validation of a prediction model and definition of new screening criteria difficult. Our precise data analyses revealed that results of this retrospective study are very robust and can therefore be expected to be valid. In order to further increase the validity of our results it would have been desirable to include more precise descriptions of ROP in terms of zone and plus disease stadium as well as type I and II ROP in the analyses (29). However, these data were not available from the SNN database.

A strength of this study is the completeness of data. Almost 7000 preterm infants below 32 weeks were evaluated over a period of 10 years. Concerning ROP, we reached data completeness of more than 97% of patients. The missing data was estimated using multivariate imputation by chained equations. Furthermore, and in contrast to many other databases the SNN database is not formed by a collaboration of certain hospitals, which might lead to a selection bias, but represents a whole population. Data of 96% of all Swiss VLBW infants born during the observation period were collected, making results representative for Switzerland. Gestational age stratified comparison with the birth registry of the federal statistical office revealed that the missing 4% of the population predominantly concern infants that died at extremely low gestational age in periphery low level neonatology units.

## Conclusion

Results of this study show that it may be possible to greatly reduce the number of infants requiring ROP screening and thereby reducing their burden and saving health care costs and resources. A prospective test of the identified model is needed before it can be applied as a general guideline.



**Statement of financial support**

The authors have no financial relationships relevant to this article to disclose.

**Disclosure statement**

The authors declare no conflict of interest.

Author accepted manuscript

**References:**

1. Hartnett ME, Penn JS 2012 Mechanisms and management of retinopathy of prematurity. *N Engl J Med* 367:2515-2526.
2. Hartnett ME 2015 Pathophysiology and mechanisms of severe retinopathy of prematurity. *Ophthalmology* 122:200-210.
3. Slidsborg C, Forman JL, Rasmussen S, *et al.* 2011 A new risk-based screening criterion for treatment-demanding retinopathy of prematurity in Denmark. *Pediatrics* 127:e598-606.
4. Darlow BA, Lui K, Kusuda S, *et al.* 2017 International variations and trends in the treatment for retinopathy of prematurity. *Br J Ophthalmol*.
5. Grupo de Trabajo Colaborativo Multicentrico para la Prevencion de la Ceguera en la Infancia por Retinopatía del P 2008 [Recommendations for Retinopathy of Prematurity screening in at-risk populations]. *Arch Argent Pediatr* 106:71-76.
6. Zin A, Florencio T, Fortes Filho JB, *et al.* 2007 [Brazilian guidelines proposal for screening and treatment of retinopathy of prematurity (ROP)]. *Arq Bras Oftalmol* 70:875-883.
7. Jefferies AL, Canadian Paediatric Society F, Newborn C 2016 Retinopathy of prematurity: An update on screening and management. *Paediatr Child Health* 21:101-108.
8. Fierson WM, American Academy of Pediatrics Section on O, American Academy of O, American Association for Pediatric O, Strabismus, American Association of Certified O 2013 Screening examination of premature infants for retinopathy of prematurity. *Pediatrics* 131:189-195.
9. Li Q, Wang Z, Wang R, Tang H, Chen H, Feng Z 2016 A Prospective Study of the Incidence of Retinopathy of Prematurity in China: Evaluation of Different Screening Criteria. *J Ophthalmol* 2016:5918736.
10. Roohipour R, Karkhaneh R, Farahani A, *et al.* 2016 Retinopathy of prematurity screening criteria in Iran: new screening guidelines. *Arch Dis Child Fetal Neonatal Ed* 101:F288-293.
11. Gerull R, Brauer V, Bassler D, *et al.* 2017 Incidence of retinopathy of prematurity (ROP) and ROP treatment in Switzerland 2006-2015: a population-based analysis. *Arch Dis Child Fetal Neonatal Ed*.
12. International Committee for the Classification of Retinopathy of P 2005 The International Classification of Retinopathy of Prematurity revisited. *Arch Ophthalmol* 123:991-999.
13. Voigt M, Rochow N, Schneider KT, *et al.* 2014 [New percentile values for the anthropometric dimensions of singleton neonates: analysis of perinatal survey data of 2007-2011 from all 16 states of Germany]. *Z Geburtshilfe Neonatol* 218:210-217.
14. Jobe AH, Bancalari E 2001 Bronchopulmonary dysplasia. *Am J Respir Crit Care Med* 163:1723-1729.
15. Bell MJ, Ternberg JL, Feigin RD, *et al.* 1978 Neonatal necrotizing enterocolitis. Therapeutic decisions based upon clinical staging. *Ann Surg* 187:1-7.
16. Papile LA, Burstein J, Burstein R, Koffler H 1978 Incidence and evolution of subependymal and intraventricular hemorrhage: a study of infants with birth weights less than 1,500 gm. *J Pediatr* 92:529-534.
17. van Buuren S G-OK 2011 Multivariate Imputation by Chained Equations in R. *Journal of Statistical Software* 45:1-67.
18. Wilson CM, Ells AL, Fielder AR 2013 The challenge of screening for retinopathy of prematurity. *Clin Perinatol* 40:241-259.
19. Wilkinson AR, Haines L, Head K, Fielder AR 2008 UK retinopathy of prematurity guideline. *Early Hum Dev* 84:71-74.
20. Nowicki M, Ostalska-Nowicka D, Kaczmarek M, Miskowiak B, Witt M 2007 The significance of VEGF-C/VEGFR-2 interaction in the neovascularization and prognosis of nephroblastoma (Wilms' tumour). *Histopathology* 50:358-364.

21. Hutchinson AK, Melia M, Yang MB, VanderVeen DK, Wilson LB, Lambert SR 2016 Clinical Models and Algorithms for the Prediction of Retinopathy of Prematurity: A Report by the American Academy of Ophthalmology. *Ophthalmology* 123:804-816.
22. Hellstrom A, Hard AL, Engstrom E, *et al.* 2009 Early weight gain predicts retinopathy in preterm infants: new, simple, efficient approach to screening. *Pediatrics* 123:e638-645.
23. Lofqvist C, Andersson E, Sigurdsson J, *et al.* 2006 Longitudinal postnatal weight and insulin-like growth factor I measurements in the prediction of retinopathy of prematurity. *Arch Ophthalmol* 124:1711-1718.
24. Eriksson L, Liden U, Lofqvist C, Hellstrom A 2014 WINROP can modify ROP screening praxis: a validation of WINROP in populations in Sormland and Vastmanland. *Br J Ophthalmol* 98:964-966.
25. Lundgren P, Stoltz Sjostrom E, Domellof M, *et al.* 2013 WINROP identifies severe retinopathy of prematurity at an early stage in a nation-based cohort of extremely preterm infants. *PLoS One* 8:e73256.
26. Choi JH, Lofqvist C, Hellstrom A, Heo H 2013 Efficacy of the screening algorithm WINROP in a Korean population of preterm infants. *JAMA Ophthalmol* 131:62-66.
27. Sun H, Kang W, Cheng X, *et al.* 2013 The use of the WINROP screening algorithm for the prediction of retinopathy of prematurity in a Chinese population. *Neonatology* 104:127-132.
28. Piyasena C, Dhaliwal C, Russell H, *et al.* 2014 Prediction of severe retinopathy of prematurity using the WINROP algorithm in a birth cohort in South East Scotland. *Arch Dis Child Fetal Neonatal Ed* 99:F29-33.
29. Good WV, Early Treatment for Retinopathy of Prematurity Cooperative G 2004 Final results of the Early Treatment for Retinopathy of Prematurity (ETROP) randomized trial. *Trans Am Ophthalmol Soc* 102:233-248; discussion 248-250.

## Acknowledgements

We would like to thank Karine Landgren-Hugentobler for proofreading and editing of the manuscript.

We also thank the following units for collaborating in this study of the SwissNeoNet: Aarau: Cantonal Hospital Aarau, Children's Clinic, Department of Neonatology (Ph. Meyer, C. Anderegg), Department of Neuropaediatrics (A. Capone Mori, D. Kaeppli); Basel: University Children's Hospital Basel, Department of Neonatology (S. Schulzke), Department of Neuropaediatrics and Developmental Medicine (P. Weber); Bellinzona: San Giovanni Hospital, Department of Pediatrics (G.P. Ramelli); Berne: University Hospital Berne, Department of Neonatology (M. Nelle), Department of Pediatrics (B. Wagner), Department of Neuropaediatrics (M. Steinlin, S. Grunt); Biel: Development and Pediatric Neurorehabilitation Center (R. Hassink); Chur: Children's Hospital Chur, Department of Neonatology (T. Riedel), Department of Neuropaediatrics (E. Keller, Ch. Killer); Fribourg: Cantonal Hospital Fribourg, Department of Neuropaediatrics (K. Fuhrer); Lausanne: University Hospital (CHUV), Department of Neonatology (J.-F. Tolsa, M. Roth-Kleiner), Department of Child Development (M. Bickle-Graz); Geneva: Department of child and adolescent, University Hospital (HUG), Neonatology Units (R. E. Pfister), Division of Development and Growth (P. S. Huppi, C. Borradori-Tolsa); Lucerne: Children's Hospital of Lucerne, Neonatal and Paediatric Intensive Care Unit (M. Stocker), Department of Neuropaediatrics (T. Schmitt-Mechelke, F. Bauder); Lugano: Regional Hospital Lugano, Department of Pediatrics (V. Pezzoli); Muensterlingen: Cantonal Hospital Muensterlingen, Department of Pediatrics (B. Erkert, A. Mueller); Neuchatel: Cantonal Hospital Neuchatel, Department of Pediatrics (M. Ecoffey); St. Gallen: Cantonal Hospital St. Gallen, Department of Neonatology (A. Malzacher), Children's Hospital St. Gallen, Neonatal and Paediatric Intensive Care Unit (J. P. Micallef), Department of Child Development (A. Lang-Dullenkopf); Winterthur: Cantonal Hospital Winterthur, Department of Neonatology (L. Hegi), Social Pediatrics Center (M. von Rhein); Zurich: University Hospital Zurich (USZ), Department of Neonatology (D. Bassler, R. Arlettaz), University Children's Hospital Zurich, Department of Neonatology (V. Bernet) and Child Development Centre (B. Latal, G. Natalucci).

# Figure legends

Figure 1: Gestational age and Birth weight of the study population. Red dots show patients with treated ROP.

Figure 2: ROC curve of c-statistic for prediction to need ROP treatment: AUC = 0.916

Gestational age (weeks)	23	24	25	26	27	28	29	30	31	Total
n (%)	19 (0.3%)	174 (2.6%)	334 (5%)	545 (8.1%)	661 (9.8%)	820 (12.2%)	1029 (15.3%)	1362 (20.3%)	1775 (26.4%)	6719 (100%)
Died >= 5 weeks GA	15.8%	4.6%	4.8%	2.2%	1.4%	1.0%	0.2%	0.5%	0.5%	1.1%
Male sex	68.4%	50.6%	56.5%	51.4%	54.5%	51.6%	53.4%	55.2%	54.6%	53.9%
Multiple births	47.4%	20.7%	21.3%	27.5%	28.0%	32.1%	36.3%	39.4%	37.9%	34.2%
Antenatal steroids	84.2%	90.2%	93.4%	90.7%	90.4%	91.4%	93.0%	92.0%	89.6%	91.2%
Caesarian section	52.6%	72.7%	79.0%	80.9%	79.3%	83.2%	80.7%	82.9%	78.0%	80.2%
SGA	5.3%	7.5%	10.2%	9.0%	11.3%	8.7%	6.3%	7.4%	8.3%	8.3%
Sepsis	26.3%	37.4%	29.0%	23.9%	17.1%	9.9%	7.3%	4.2%	3.3%	10.2%
NEC	15.8%	9.2%	2.4%	3.3%	2.6%	2.6%	1.7%	1.5%	0.8%	2.0%
Severe IVH	15.8%	9.8%	13.2%	7.0%	6.8%	2.9%	3.1%	2.6%	1.5%	3.9%
O <sub>2</sub> days	65.2 (27.6)	69.2 (40.1)	57 (38.4)	42.6 (35.3)	34.4 (30.0)	22.4 (27.5)	12.4 (18.3)	7.0 (12.8)	4.3 (15.3)	18.8 (29.1)
BPD	50.0%	45.5%	32.7%	23.8%	17.0%	10.9%	6.5%	3.6%	2.2%	10.0%
Length of stay (days)	126.6 (67.4)	120.8 (39.9)	110.3 (38.2)	95.5 (31.3)	85.7 (27.1)	73.8 (30.8)	62.2 (22.9)	51.5 (20.3)	42.6 (17.6)	65.4 (33.6)
ROP stage 0	62.5%	56.0%	61.5%	76.9%	86.3%	91.7%	95.5%	97.7%	98.6%	90.5%

<b>ROP stage 1</b>	31.2%	9.6%	12.6%	11.3%	8.7%	5.4%	2.6%	1.6%	1.1%	4.7%
<b>ROP stage 2</b>	0.0%	13.9%	16.6%	7.2%	3.7%	2.2%	1.0%	0.5%	0.2%	2.9%
<b>ROP stage 3</b>	6.2%	20.5%	9.2%	4.3%	1.2%	0.5%	0.7%	0.2%	0.1%	1.8%
<b>ROP stage 4</b>	0.0%	0.0%	0.0%	0.0%	0.0%	0.1%	0.0%	0.0%	0.0%	0.0%
<b>ROP stage 5</b>	0.0%	0.0%	0.0%	0.2%	0.0%	0.1%	0.1%	0.0%	0.0%	0.0%
<b>Severe ROP</b>	6.2%	20.5%	9.2%	4.5%	1.2%	0.8%	0.8%	0.2%	0.1%	1.9%
<b>ROP intervention</b>	5.3%	14.5%	7.3%	2.6%	1.1%	0.5%	0.0%	0.2%	0.1%	1.2%
<b>ROP intervention anti-VEGF</b>	0.0%	0.6%	1.2%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.1%
<b>ROP intervention lasertherapy</b>	5.3%	13.9%	6.1%	2.6%	1.1%	0.5%	0.0%	0.2%	0.1%	1.1%

Table 1 Patient characteristics stratified by gestational age

GA: gestational age; SGA: small for gestational age (<10th percentile) NEC: necrotizing enterocolitis; IVH: intraventricular hemorrhage; BPD: bronchopulmonary dysplasia; ROP retinopathy of prematurity

GA	Birth weight	Birth weight z-score	Gender	Multiple birth	Surfactant	NEC	IVH Grade	Days O <sub>2</sub>	Days CPAP	Days mech. ventilation	Days hospitalization
27+0	1140g	0.576	M	yes	no	no	0	72	32	0	82
27+1	800g	-0.908	M	yes	yes	no	2	69	37	24	81
27+1	700g	-1.073	F	yes	yes	no	3	68	39	5	81
27+1	480g	-2.306	M	no	yes	yes	0	62	55	1	113
27+2	960g	0.109	F	no	no	no	0	12	33	0	62
27+4	462g	-2.155	F	no	no	no	0	102	69	11	102
28+0	770g	-1.345	M	no	no	no	2	111	37	0	116
28+1	517g	-2.034	F	yes	yes	no	2	180	70	38	186
28+3	680g	-1.412	F	no	no	no	0	52	39	2	84
28+4	660g	-1.489	F	yes	no	no	0	124	69	17	215
30+1	1530g	0.503	F	no	yes	no	0	10	0	10	52
30+5	1370g	0.006	F	no	yes	no	0	36	9	5	72

31+4	1300g	-0.725	M	no	no	no	0	1	0	0	48
------	-------	--------	---	----	----	----	---	---	---	---	----

Table 2: Parameters of patients with a gestational age above 26 weeks needing ROP treatment  
GA: gestational age; UA pH: pH in umbilical artery; NEC: necrotizing enterocolitis; IVH: intraventricular hemorrhage

	est	SE	p-value	Odds ratios	95% CI
<b>(Intercept)</b>	-1.257	0.653	0.054	NA	NA
<b>GA (weeks/days)</b>	-0.75	0.109	0	0.472	0.38-0.59
<b>SGA (yes)</b>	0.666	0.389	0.087	1.946	0.91-4.17
<b>Multiple birth (yes)</b>	0.669	0.313	0.033	1.952	1.06-3.61
<b>Surfactant administration (yes)</b>	-0.725	0.336	0.031	0.484	0.25-0.94
<b>O2 (day)</b>	0.013	0.003	0	1.013	1.01-1.02
<b>CPAP (day)</b>	0.012	0.004	0.008	1.012	1.00-1.02
<b>Ventilation (day)</b>	0.015	0.007	0.037	1.015	1.00-1.03


Table 3: Patient characteristics used in the final prediction model

est: maximum likelihood estimator used for each parameter to calculate probability using formula above ( $b_k$ ); SE: standard error; OR: odds ratio; lo 95 / hi 95: 95% confidence interval.

cutoff at predicted value	n needed to test	True Pos	False Neg	Sensitivity	Specificity
0.000	2108	20	0	1.000	0.000
0.001	1452	19	1	0.950	0.314
0.002	1102	19	1	0.950	0.481
0.040	284	19	1	0.950	0.873
0.041	278	19	1	0.950	0.876
0.042	272	18	2	0.900	0.878
0.045	261	18	2	0.900	0.884
0.046	255	17	3	0.850	0.886
0.054	234	17	3	0.850	0.896
0.055	231	16	4	0.800	0.897
0.068	184	16	4	0.800	0.920
0.069	178	15	5	0.750	0.922
0.071	177	15	5	0.750	0.922
0.072	174	14	6	0.700	0.923
0.073	171	13	7	0.650	0.924
...					

Table 4: Cutoff values for predicted ROP intervention probability.





# ROC Rop Intervention Model

